Author Index

ABADIE ANNIE & PROUVOST-DANON ANNIE Specific and total IgE responses to antigenic stimuli in Brown–Norway, Lewis and Sprague–Dawley rats, 559
ALMERIGOGNA F., *see* ROMAGNANI S.
ANDRÉ C.M., *see* LEWIS J.G.
ANDREW E., *see* REYNOLDS J.
ARIMORI S., *see* HIRAMATSU KAZUKO
ASHERSON G.L., *see* THOMAS W.R.
ASKENASE P.W., *see* OGILVIE BRIDGET M.

BABIUK L.A., *see* HO C.K.
BALTZ MARILYN L., *see* PEPYS M.B.
BILLING R., *see* WEISBART R.H.
BITO Y., *see* HIROTA Y.
BLOCH K.J. *see* LAKE A.M.
BLOMBERG MARY VON & SCHEPER R.J. Hapten-cell conjugates in DNCB contact sensitivity. *In vitro* stimulation with DNP-conjugates optimally inducing contact sensitivity *in vivo*, 291
BOLDT D.H. & LYONS RUTH D. Fractionation of human lymphocytes with plant lectins. III. Identification of cells regulating the *in vitro* response to L-phytohaemagglutinin, 517
BRANDTZAEG P., *see* KORSRUD F.R.
BRASH A.R., *see* GOETZEL E.J.
BROOKS C.G. & FLANNERY G.R. Quantitative studies of natural immunity to solid tumours in rats. Persistence of natural immunity throughout reproductive life, and absence of suppressor cells in infant rats, 187
BROWN SUSAN, *see* KEMP A.S.
BULLEN J.J., LEACH P. ANN & LANGE LINDA Abolition of resistance to *Klebsiella pneumoniae* by anti-polymorphonuclear leucocyte IgG, 47
BUSKJAER L., *see* MORTENSEN J.P.

CASTES B. MARIANELLA, *see* ORBACH-ARBOUYS SIMONE
CATALONA W.J., RATLIFF T.L. & MCCOOL R.E. Characterization of cell-mediated cytolytic mechanism against human transitional cell carcinoma line, 253J, 119
CHAOUAT G. & VOISIN G.A. Regulatory T-cell subpopulations in pregnancy. II. Evidence for suppressive activity of the late phase of MLR, 239
CHEDID L., *see* LÖWY ILANA
COLLINS F.M., *see* WATSON SUSAN R.
COMOGLIO P.M., *see* PRAT MAGA
COOK R.M. Spontaneous cytotoxicity of murine cells treated with the interferon inducers BRL 5907 and BRL 10739, 151
COOMBS R.R.A., *see* WILSON ANNE B.
CORNWELL C.G. III, MICHAELSEN T.E. & SANDERS W.H. Comparative studies of a monoclonal IgG_λ and two IgE_κ components from an individual patient: evidence for shared idiotypic determinants between the two IgE proteins but not with the IgG_λ protein, 509
COTTRELL B.J., STURROCK R.F. & VAN HOEGAERDEN M. An immunosuppressive factor in the serum of baboons (*Papio anubis*) infected with *Schistosoma mansoni*, 587

DALLMAN MARGARET, *see* NAYSMITH J.D.
DAVIES A.J.S., *see* HUCHET R.
DE BRACCO MARÍA M. DE E., *see* RIMOLDI MARÍA THERESA
DESAYMARD CATHERINE, SHINTON R.A. & WALDMANN H. Differences in calcium requirements for B-cell tolerance and immunity, 61
DOENHOFF M.J., *see* PEPYS M.B.

EDWARDS A.J., *see* THOMAS W.R.
ELSON C.J., *see* NAYSMITH J.D.
ENDOH M., *see* HIRAMATSU KAZUKO

FAUCI A.S., *see* STEVENSON H.C.
FAUCI S., *see* KATZ P.
FLANNERY G.R., *see* BROOKS C.G.
FLETCHER ELAIN, *see* NAYSMITH J.D.
FORD P.M. & KOSATKA IVA. The effect of *in situ* formation of antigen–antibody complexes in the glomerulus on subsequent glomerular localization of passively administered immune complexes, 337
FORSBERG J.-G., *see* KALLAND T.
FRANCIS DAPHNE H., *see* TAYLOR W.A.

GERDES J., KLATT UTE & STEIN H. Xenoantiserum to human C3 receptors: its preparation and effect on the C3b and C3d receptors of tonsil cells and the C3b receptors of erythrocytes and neutrophils, 75
GILL T.J. III, *see* NOEMAN S.A.
GIUDIZI GRAZIA M., *see* ROMAGNANI S.
GOETZL E.J., BRASH A.R., TAUBER A.I., OATES J.A. & HUBBARD W.C. Modulation of human neutrophil function by monohydroxy-eicosatetraenoic acids, 489
GOLDSCHNEIDER I., *see* RITTER MARY A.
GOLSTEIN P., *see* MACLENNAN I.C.M.
GOREN RIVKA & NELKEN D. Normal immunosuppressive protein. Isolation of a glycoprotein active fraction, 305
GOTCH FRANCES M., *see* MACLENNAN I.C.M.
GRANDJON D., *see* HUCHET R.
GRIFFIN F.M. JR, *see* SHAW DENISE R.
GUÉNET J.-L., *see* MARCHAL G.
GYURE L., *see* REYNOLDS J.

HALL J.G., HOPKINS J. & REYNOLDS J. Studies of efferent lymph cells from nodes stimulated with oxazolone, 141
HALL J.G., *see* REYNOLDS J.
HAMILTON FIONA, *see* WARDLEY R.C.
HAMURO J., RÖLLINGHOFF M. & WAGNER H. Induction of cytotoxic peritoneal exudate cells by T-cell immune adjuvants of the $\beta(1\rightarrow3)$ glucan-type lentinan and its analogues, 549

HANAOKA M., *see* TAKABAYASHI A.
HARD R.C. JR Sequential changes in serum immunoglobulin levels in young RFM mice with host-versus-graft disease, 453
HARRIS G. The effect of ethidium bromide on the response of spleen explants of primed rabbits to sheep red cells (SRC), 165
HEATLEY R.V., *see* STARK J.M.
HENDERSON D.C., PARKER DARIEN & TURK J.L. Dissociation between tumour resistance and delayed-type hypersensitivity to tumour-associated antigens in the mouse, 1
HICKMAN SUZANNE E., *see* SHAW DENISE R.
HIRAMATSU KAZUKO, SAKAI H., ENDOH M. & ARIMORI S. Surface properties of LDL-binding lymphocytes in human peripheral blood, 311
HIROTA Y., SUZUKI T. & BITO Y. The development of unusual B-cell functions in the testosterone-propionate-treated chicken, 29
HIROTA Y., SUZUKI T. & BITO Y. The B-cell development independent of the bursa of Fabricius but dependent upon the thymus in chickens treated with testosterone propionate, 37
HO C.K. & BABIUK L.A. Immune mechanisms against canine distemper. III. Role of complement lysis in the immunity and persistent infection of canine distemper virus, 231
HOLLIMAN A., *see* TAUSSIG M.J.
HOPKINS J., *see* HALL J.G.
HUBBARD W.C., *see* GOETZL E.J.
HUCHET R., OLSSON L., GRANDJON O. & DAVIES A.J.S. Hapten-specific unresponsiveness in mice. II. Fate and distribution of ^{14}C-TNBSA injected at tolerogenic doses, 275

INMAN J.K., *see* SNIPPE H.

JASANI B. & STARK J.M. Endotoxin-induced appearance of peroxidase-positive cells in the white pulp of mouse spleen, 533
JOHANNESEN L., *see* SNIPPE H.

KALLAND T. & FORSBERG J.-G. Permanent inhibition of capping of spleen lymphocytes from neonatally oestrogen-treated female mice, 281
KATZ P. & FAUCI S. Antibody-dependent cellular cytotoxicity mediated by subpopulations of human T lymphocytes: killing of human erythrocytes and autologous lymphoid cells, 407
KEMP A.S. & BROWN SUSAN Inhibition of neutrophil migration by aggregated immunoglobulin attached to micropore membranes, 527
KLATT UTE, *see* GERDES J.
KORSRUD F.R. & BRANDTZAEG P. Quantitative immunohistochemistry of immunoglobulin- and J-chain-producing cells in human parotid and submandibular salivary glands, 129
KOSATKA IVA, *see* FORD P.M.
KOYAMA K., *see* TANIGAKI N.
KRAKOWKA S. Mechanisms of E-rosette formation by mitogen-stimulated canine lymphocytes, 255

LACHMANN P.J., *see* WILSON ANNE B.
LAKE A.M., BLOCH K.J., SINCLAIR K.J. & WALKER W.A. Anaphylactic release of intestinal goblet cell mucus, 173
LAMM L.U., *see* MORTENSEN J.P.
LAMON E.W., *see* SHAW DENISE R.
LANGE LINDA, *see* BULLEN J.J.
LAWMAN M.J., *see* WARDLEY R.C.
LEACH P. ANN, *see* BULLEN J.J.
LECLERC C., *see* LÖWY ILANA
LEWIS J.G. & ANDRÉ C.M. Effect of human α_2HS glycoprotein on mouse macrophage function, 317
LIZZIO ELAINE, *see* SNIPPE H.
LOCKE JOAN, *see* STARK J.M.
LÖWY ILANA, LECLERC C. & CHEDID L. Induction of antibodies directed against self and altered-self determinants by a synthetic adjuvant, muramyl dipeptide and some of its derivatives, 441
LYONS RUTH D., *see* BOLDT D.H.

MACLENNAN I.C.M., GOTCH FRANCES M. & GOLSTEIN P. Limited specific T-cell mediated cytolysis in the absence of extracellular Ca^{2+}, 109
MCCOOL R.E., *see* CATALONA W.J.
MCLEOD RIMA & REMINGTON J.S. Inhibition or killing of an intracellular pathogen by activated macrophages is abrogated by TLCK or aminophylline, 597
MAGRO A.M. Creatine phosphokinase in rat mast cells, 323
MAN D.A., *see* MINTA J.O.
MARCHAL G., MILON GENEVIÈVE & GUÉNET J.-L. Altered circulation of lymphocytes mediating delayed-type hypersensitivity when primed in W^f/W^f mice, 269
MASUDA T., *see* TAKABAYASHI A.
MATSUMOTO S., *see* TONO-OKA T.
MERCHANT B., *see* SNIPPE H.
MICHAELSEN T.E., *see* CORNWELL C.G. III
MILON GENEVIÈVE, *see* MARCHAL G.
MINTA J.O. & MAN D.A. Immunological, structural and functional relationships between an anti-complementary protein from *Crotalus atrox* venom, cobra venom factor and human C3, 501
MISRA D.N., *see* NOEMAN S.A.
MITCHELL G.F., RAJASEKARIAH G.R. & RICKARD M.D. A mechanism to account for mouse strain variation in resistance to the larval cestode, *Taenia taeniaeformis*, 479
MOORE M., *see* POTTER M.R.
MORRIS R.J., *see* RITTER MARY A.
MORTENSEN J.P., BUSKJAER L. & LAMM L.U. Studies on the C2-deficiency gene in man, 539
MUSALLAM R., *see* PEPYS M.B.

NAKAYAMA M., *see* TONO-OKA T.
NAYSMITH J.D., ELSON C.J., DALLMAN MARGARET, FLETCHER ELAIN & ORTEGA-PIERRES MARIA G. Anti-erythrocyte autoantibody production in mice associated with the injection of rat erythrocytes. I. Regulation of the response by suppressor cells, 409
NICOLETTI P.L., *see* ROMAGNANI S.
NOEMAN S.A., MISRA D.N. & GILL T.J. III Characterization of the T antigen and T agglutinin in inbred rats, 219

OATES J.A., *see* GOETZL E.J.
OGILVIE BRIDGET M., ASKENASE P.W. & ROSE M. ELAINE Basophils and eosinophils in three strains of rats and in athymic (nude) rats following infection with the nematodes *Nippostrongylus brasiliensis* or *Trichinella spiralis*, 385

OLSSON L., *see* HUTCHET R.
ORBACH-ARBOUYS SIMONE & CASTES B. MARIANELLA Suppression of T-cell responses to histocompatibility antigens by BCG pre-treatment, 263
ORTEGA-PIERRES MARIA G., *see* NAYSMITH J.D.

PARKER DARIEN, *see* HENDERSON D.C.
PAWELEC G. The induction of suppressor cells in mixed leucocyte cultures and in mixed leucocyte–non-lymphoid cell cultures, 11
PEPYS M.B., BALTZ MARILYN L., MUSALLAM R. & DOENHOFF M.J. Serum protein concentrations during *Schistosoma mansoni* infection in intact and T-cell deprived mice. I. The acute phase proteins, C3 and serum amyloid P-component (SAP), 249
POTTER M.R. & MOORE M. The effect of BCG stimulation on natural cytotoxicity in the rat, 427
PRAT MARIA, TATO F., TARONE G. & COMOGLIO P.M. Target antigens for antibodies and complement at the cell surface of RSV-transformed fibroblasts, 179
PRESSMAN D., *see* TANIGAKI N.
PRICHARD-THOMAS SARAH, *see* WILSON ANNE B.
PROUVOST-DANON ANNIE, *see* ABADIE ANNIE

RABINOWITZ RUTH & SCHLESINGER M. Inhibition of the activity of cytotoxic murine T lymphocytes by antibodies to idiotypic determinants, 93
RAJASEKARIAH G.R., *see* MITCHELL G.F.
RATLIFF T.L., *see* CATALONA W.J.
REYNOLDS J., GYURE L., ANDREW E. & HALL J.G. Studies of transport of polyclonal IgA antibody from blood to bile in rats, 463
REYNOLDS J., *see* HALL J.C.
RICCI M., *see* ROMAGNANI S.
RICKARD M.D., *see* MITCHELL G.F.
RIMOLDI MARÍA TERESA & DE BRACCO MARÍA M. DE E. *In vitro* inactivation of complement by a serum factor present in Junin-virus infected guinea-pigs, 159
RITTER MARY A., MORRIS R.J. & GOLDSCHNEIDER I. Hidden Thy-1 antigen in a subpopulation of mouse bone marrow cells, 375
RITTER MARY A. & MORRIS R.J. Thy-1 antigen: selective association in lymphoid organs with the vascular basement membrane involved in lymphocyte recirculation, 85
ROETERINK C.H., *see* VAN ROOIJEN N.
REMINGTON J.S., *see* MCLEOD RIMA
RÖLLINGHOFF M., *see* HAMURO J.
ROMAGNANI S., GIUDIZI GRAZIA M., ALMERIGOGNA F., NICOLETTI P.L. & RICCI M. Protein A of *Staphylococcus aureus* is mitogenic for IgG-bearing, but also for a subpopulation of IgM- and/or IgD-bearing human lymphocytes, 417
ROSE ELAINE M., *see* OGILVIE BRIDGET M.
RUDNICKA W. Suppressor activity of blood lymphocytes from guinea-pigs treated with ALS, 435

SAKAI H., *see* HIRAMATSU KAZUKO
SAKANO T. Enhancement of T-cell response to PWM by neuraminidase-treated non-T cells, 203
SANDERS W.H., *see* CORNWELL C.G. III
SCHEPER R.J., *see* BLOMBERG MARY VON
SCHLESINGER M., *see* RABINOWITZ RUTH
SCULLY C. M. Comparative opsonic activity for *Streptococcus mutans* in oral fluids, and phagocytic activity of blood, crevicular, and salivary polymorphonuclear leucocytes in rhesus monkeys, 101
SHALEV A. Pregnancy-induced H-Y antibodies and their transmission to the foetus in rats, 285
SHAW DENISE R., SHAW M.W., HICKMAN SUZANNE E., LAMON E.W. & GRIFFIN F.M. JR Sodium azide inhibition of complement-mediated functions, 53
SHAW M.W., *see* SHAW DENISE R.
SHELDON DIANNE, *see* TAYLOR W.A.
SHINTON R.A., *see* DESAYMARD CATHERINE
SINCLAIR K.J., *see* LAKE A.M.
SNIPPE H., WILLERS J.M.N., INMAN J.K. & MERCHANT B. The specificity of antibody formation in mice following immunization with hapten–carrier complexes mixed with the surfactant, dimethyl dioctadecyl ammonium bromide, 361
SNIPPE H., JOHANNESEN L., LIZZIO ELAINE & MERCHANT B. Variable expression of delayed hypersensitivity in different mouse strains using dimethyl dioctadecyl ammonium bromide as adjuvant, 399
STARK J.M., LOCKE JOAN & HEATLEY R.V. Immunogenicity of lipid-conjugated antigens. I. The influence of chain length and degree of conjugation on induction of antibody in mice, 345
STARK J.M., MATTHEWS N. & LOCKE JOAN Immunogenicity of lipid-conjugated antigens. II. Anti-complementary activity and antigen trapping in the spleen, 353
STARK J.M., *see* JASANI B.
STEIN H., *see* GERDES J.
STEVENSON H.C. & FAUCI A.S. Activation of human B lymphocytes. XII. Differential effects of *in vitro* cyclophosphamide on human lymphocyte subpopulations involved in B-cell activation, 391
STURROCK R.F., *see* COTTRELL B.J.
SUZUKI T., *see* HIROTA Y.
SUZUKI S., *see* TAKABAYASHI A.

TAKABAYASHI A., MASUDA T., SUZUKI S. & HANAOKA M. Immunological properties of Fc receptors on lymphocytes. VIII. The behaviour of FcR^+ and FcR^- T cells in cell-mediated immune responses, 209
TANIGAKI N., TOSI R., KOYAMA K. & PRESSMAN D. Purification and separation of subsets of human Ia molecules by papain digestion, 613
TARONE G., *see* PRAT MARIA
TATO F., *see* PRAT MARIA
TAUBER A.I., *see* GOETZL E.J.
TAUSSIG M.J., HOLLIMAN A. & WRIGHT L.J. Hybridization between T and B lymphoma cell lines, 57
TAYLOR W.A., SHELDON DIANNE & FRANCIS DAPHNE H. IgE antibody formation in BALB/c mice without adjuvant: induction of responses to grass pollen extract and to a hapten–carrier conjugate, 581
THOMAS W.R., EDWARDS A.J., WATKINS MADELEINE C. & ASHERSON G.L. Distribution of immunogenic cells after painting with the contact sensitizers fluorescein isothiocyanate and oxazolone. Different sensitizers form immunogenic complexes with different cell populations, 21
THOMAS W.R., VARDINON NURIT, WATKINS MADELEINE C. & ASHERSON G.L. Antigen-specific mast cell degranulation in contact sensitivity to picryl chloride. An early event, 331

TONO-OKA T., NAKAYAMA M. & MATSUMOTO S. Human monocyte-derived chemotactic factor for granulocytes, 605
TOSI R., *see* TANIGAKI N.
TURK J.L., *see* HENDERSON D.C.

VAN HOEGAERDEN M., *see* COTTRELL B.J.
VAN ROOIJEN N. & ROETERINK C.H. Phagocytosis and lymphocyte migration: evidence that lymphocyte trapping in the spleen following carbon injection is not due to direct lymphocyte–macrophage adherence, 569
VARDINON NURIT, *see* THOMAS W.R.
VOISIN G.A., *see* CHAOUAT G.

WAGNER H., *see* HAMURO J.
WALDMANN H., *see* DESAYMARD CATHERINE
WALKER W.A., *see* LAKE A.M.
WARDLEY R.C., LAWMAN M.J. & HAMILTON FIONA The establishment of continuous macrophage cell lines from peripheral blood monocytes, 67
WATKINS MADELEINE C., *see* THOMAS W.R.
WATSON SUSAN R. & COLLINS F.M. Development of suppressor T cells in mice heavily infected with mycobacteria, 367
WEISBART R.H. & BILLING R. Requirement for B-lymphocyte alloantigen in the production of neutrophil migration inhibition factor from T lymphocytes (NIF-T), 575
WILLERS J.M.N., *see* SNIPPE H.
WILSON ANNE B., PRICHARD-THOMAS SARAH, LACHMANN P.J. & COOMBS R.R.A Receptors on guinea-pig erythrocytes specific for cell-bound fourth component of human complement (C4), 195
WRIGHT L.J., *see* TAUSSIG M.J.

Subject Index

Anaphylactic release of intestinal goblet cell mucus, 173
Antibodies, induction of, directed against self and altered-self determinants by a synthetic adjuvant, muramyl dipeptide and some of its derivatives, 441
Autoantibody production in mice, anti-erythrocyte, associated with the injection of rat erythrocytes. I. Regulation of the response by suppressor cells, 469

BCG pre-treatment, suppression of T-cell responses to histocompatibility antigens by, 263
BCG stimulation, the effect of, on natural cytotoxicity in the rat, 427
B-cell development, the, independent of the bursa of Fabricius but dependent upon the thymus in chickens treated with testosterone propionate, 37
B-cell functions, the development of unusual, in the testosterone-propionate-treated chicken, 29
B-cell tolerance and immunity, differences in calcium requirements for, 61
B-lymphocyte alloantigen, requirement for, in the production of neutrophil migration inhibition factor from T lymphocytes (NIF-T), 575
B lymphocytes., Activation of human, XII. Differential effects of *in vitro* cyclophosphamide on human lymphocyte subpopulations involved in B-cell activation, 391
B lymphoma cell lines, hybridization between T and, 57
Basinophils and eosinophils in three strains of rats and in athymic (nude) rats following infection with the nematodes *Nippostrongylus brasiliensis* or *Trichinella spiralis*, 385
Bone marrow cells, hidden Thy-1 antigen in a subpopulation of mouse, 375

C2-deficiency gene in man, studies on, 539
C3, human, immunological, structural and functional relationships between an anti-complementary protein from *Crotalus atrox* venom, cobra venom factor and, 501
C3 receptors: Xenoantiserum to human, its preparation and effect on the C3b and C3d receptors of tonsil cells and the C3b receptors of erythrocytes and neutrophils, 75
Calcium requirements for B-cell tolerance and immunity, differences in, 61
Carcinoma line, 253J, characterization of cell-mediated cytolytic mechanisms against human transitional cell, 119
Cobra venom factor and human C3, immunological, structural and functional relationships between an anti-complementary protein from *Crotalus atrox* venom, 501
Complement (C4), human, receptors on guinea-pig erythrocytes specific for cell-bound fourth component of, 195
Complement, *in vitro* inactivation of, by a serum factor present in Junin-virus infected guinea-pigs, 159
Complement-mediated functions, sodium azide inhibition of, 53
Contact sensitivity, Hapten–cell conjugates in DNCB, *In vitro* stimulation with DNP-conjugates optimally inducing contact sensitivity *in vivo*, 291
Contact sensitivity to picryl chloride. An early event, antigen-specific mast cell degranulation in, 331
Creatine phosphokinase in rat mast cells, 323
Crotalus atrox venom, cobra venom factor and human C3, immunological, structural and functional relationships between an anti-complementary protein from, 501
Cyclophosphamide, activation of human B lymphocytes. XII. Differential effects of *in vitro*, on human lymphocyte subpopulations involved in B-cell activation, 391
Cytolysis, limited specific T-cell mediated, in the absence of extracellular Ca^{2+}, 109
Cytolytic mechanisms, characterization of cell-mediated, against human transitional cell carcinoma line, 253J, 119
Cytotoxic murine T lymphocytes, inhibition of the activity of, by antibodies to idiotypic determinants, 93
Cytotoxicity, antibody-dependent cellular, mediated by subpopulations of human T lymphocytes: killing of human erythrocytes and autologous lymphoid cells, 407
Cytotoxicity in the rat, the effect of BCG stimulation on natural, 427
Cytotoxicity, spontaneous, of murine cells treated with the interferon inducers BRL 5907 and BRL 10739, 151

Delayed hypersensitivity, variable expression of, in different mouse strains using dimethyl dioctadecyl ammonium bromide as adjuvant, 399
Delayed-type hypersensitivity, dissociation between tumour resistance and, to tumour-associated antigens in the mouse, 1
Delayed-type hypersensitivity, altered circulation of lymphocytes mediating, when primed in W^f/W^f mice, 269
Dimethyl dioctadecyl ammonium bromide, the specificity of antibody formation in mice following immunization with hapten–carrier complexes mixed with the surfactant, 361
Dimethyl dioctadecyl ammonium bromide as adjuvant, variable expression of delayed hypersensitivity in different mouse strains using, 399
Distemper, Immune mechanisms against canine, III. Role of complement lysis in the immunity and persistent infection of canine distemper virus, 231

E-rosette formation, mechanisms of, by mitogen-stimulated canine lymphocytes, 255
Endotoxin-induced appearance of peroxidase-positive cells in the white pulp of the mouse spleen, 533
Eosinophils, basinophils and, in three strains of rats and in athymic (nude) rats following infection with the nematodes *Nippostrongylus brasiliensis* or *Trichinella spiralis*, 385
Erythrocytes, receptors on guinea-pig, specific for cell-bound fourth component of human complement (C4), 195

Ethidium bromide, the effect of, on the response of spleen explants of primed rabbits to sheep red cells (SRC), 165

Fc receptor, immunological properties of, on lymphocytes. VIII. The behaviour of FcR^+ and FcR^- T cells in cell-mediated immune responses, 209
Fibroblasts, target antigens for antibodies and complement at the cell surface of RSV-transformed, 179
Fluorescein isothiocyanate and oxazolone, distribution of immunogenic cells after painting with the contact sensitizers, Different sensitizers form immunogenic complexes with different cell populations, 21

Glomerulus, the effect of *in situ* formation of antigen-antibody complexes in the, on subsequent glomerular localization of passively administered immune complexes, 337
Glycoprotein active fraction, normal immunosuppressive protein. Isolation of a, 305
Glycoprotein, effect of human α_2HS, on mouse macrophage function, 317
Goblet cell mucus, anaphylactic release of intestinal, 173
Granulocytes, human monocyte-derived chemotactic factor for, 605

H-Y antibodies, pregnancy-induced, and their transmission to the foetus in rats, 285
Hapten–cell conjugates in DNCB contact sensitivity. *In vitro* stimulation with DNP-conjugates optimally inducing contact sensitivity *in vivo*, 291
Hapten-specific unresponsiveness in mice. II. Fate and distribution of ^{14}C-TNBSA injected at tolerogenic doses, 275
Host-versus-graft disease, sequential changes in serum immunoglobulin levels in young RFM mice with, 453

Ia molecules, purification and separation of subsets of human, by papain digestion, 613
IgA antibody, studies of transport of polyclonal, from blood to bile in rats, 463
IgE antibody formation in BALB/c mice without adjuvant: induction of responses to grass pollen extract and to a hapten–carrier conjugate, 581
IgE responses, specific and total, to antigenic stimuli in Brown–Norway, Lewis and Sprague–Dawley rats, 559
IgE_K components, comparative studies of a monoclonal IgG_λ and two, from an individual patient: evidence for shared idiotypic determinants between the two IgE proteins but not with the IgG_λ protein, 509
IgG, abolition of resistance to *Klebsiella pneumoniae* by anti-polymorphonuclear leucocyte, 47
IgG_λ, comparative studies of a monoclonal, and two IgE_K components from an individual patient: evidence for shared idiotypic determinants between the two IgE proteins but not with the IgG_λ protein, 509
Immunoglobulin, aggregated, inhibition of neutrophil migration by, attached to micropore membranes, 527
Immunoglobulin- and J-chain-producing cells, quantitative immunohistochemistry of, in human parotid and submandibular salivary glands, 129
Immunoglobulin levels, serum, sequential changes in, in young RFM mice with host-versus-graft disease, 453
Interferon inducers BRL 5907 and BRL 10739, spontaneous cytotoxicity of murine cells treated with the, 151

J-chain-producing cells, quantitative immunohistochemistry of immunoglobulin- and, in human parotid and submandibular salivary glands, 129
Junin-virus infected guinea-pigs, *in vitro* inactivation of complement by a serum factor present in, 159

Klebsiella pneumoniae, abolition of resistance to, by anti-polymorphonuclear leucocyte IgG, 47

Leucocyte cultures, the induction of suppressor cells in mixed, and in mixed leucocyte–non-lymphoid cell cultures, 11
Lipid-conjugated antigens., Immunogenicity of, I. The influence of chain length and degree of conjugation on induction of antibody in mice, 345
Lipid-conjugated antigens., Immunogenicity of, II. Anti-complementary activity and antigen trapping in the spleen, 353
Lymph cells, studies of efferent, from nodes stimulated with oxazolone, 141
Lymphocyte migration:, phagocytosis and, evidence that lymphocyte trapping in the spleen following carbon injection is not due to direct lymphocyte–macrophage adherence, 569
Lymphocyte recirculation, Thy-1 antigen: selective association in lymphoid organs with the vascular basement membrane involved in, 85
Lymphocytes, altered circulation of mediating delayed-type hypersensitivity when primed in w^f/w^f mice, 269
Lymphocytes, fractionation of human, with plant lectins. III. Identification of cells regulating the *in vitro* response to L-phytohaemagglutinin, 517
Lymphocytes, inhibition of the activity of cytotoxic murine T, by antibodies to idiotypic determinants, 93
Lymphocytes, mechanisms of E-rosette formation by mitogen-stimulated canine, 255
Lymphocytes, permanent inhibition of capping of spleen, from neonatally oestrogen-treated female mice, 281
Lymphocytes, suppressor activity of blood, from guinea-pigs treated with ALS, 435
Lymphocytes, surface properties of LDL-binding, in human peripheral blood, 311
Lymphoma cell lines, hybridization between T and B, 57

Macrophage cell lines, the establishment of continuous, from peripheral blood monocytes, 67
Mast cells, creatine phosphokinase in rat, 323
Monocytes, the establishment of continuous macrophage cell lines from peripheral blood, 67
Muramyl dipeptide, induction of antibodies directed against self and altered-self determinants by a synthetic adjuvant, and some of its derivatives, 441
Mycobacteria, development of suppressor T cells in mice heavily infected with, 367

Neuraminidase-treated non-T cells, enhancement of T-cell response to PWM by, 203
Neutrophil function, modulation of, by monohydroxy-eicosatetraenoic acids, 489
Neutrophil migration, inhibition of, by aggregated immunoglobulin attached to micropore membranes, 527

Neutrophil migration inhibition factor, requirement for B-lymphocyte alloantigen in the production of, from T lymphocytes (NIF-T), 575

Oestrogen-treated female mice, permanent inhibition of capping of spleen lymphocytes from neonatally, 281
Opsonic activity, comparative, for *Streptococcus mutans* in oral fluids, and phagocytic activity of blood, crevicular, and salivary polymorphonuclear leucocytes in rhesus monkeys, 101
Oxazolone, distribution of immunogenic cells after painting with the contact sensitizers fluorescein isothiocyanate and, Different sensitizers form immunogenic complexes with different cell populations, 21
Oxazolone, studies of efferent lymph cells from nodes stimulated with, 141

Papain digestion, purification and separation of subsets of human Ia molecules by, 613
Pathogen, inhibition or killing of an intracellular pathogen by activated macrophages is abrogated by TLCK or aminophylline, 597
Peritoneal exudate cells, induction of cytotoxic, by T-cell immune adjuvants of the $\beta(1\rightarrow3)$ glucan-type lentinan and its analogues, 549
Peroxidase-positive cells, endotoxin-induced appearance of, in the white pulp of the mouse spleen, 533
Phagocytosis and lymphocyte migration: evidence that lymphocyte trapping in the spleen following carbon injection is not due to direct lymphocyte–macrophage adherence, 569
L-Phytohaemagglutinin, fractionation of human lymphocytes with plant lectins. III. Identification of cells regulating the *in vitro* response to, 517
Picryl chloride. An early event, antigen-specific mast cell degranulation in contact sensitivity, 331
Pregnancy-induced H-Y antibodies and their transmission to the foetus in rats, 285
Protein A of *Staphylococcus aureus* is mitogenic for IgG-bearing, but also for a subpopulation of IgM- and/or IgD-bearing human lymphocytes, 417
PWM, enhancement of T-cell response to, by neuraminidase-treated non-T cells, 203

RSV-transformed fibroblasts, target antigens for antibodies and complement at the cell surface of, 179

Schistosoma mansoni, an immunosuppressive factor in the serum of baboons (*Papio anubis*) infected with, 587
Schistosoma mansoni infection, serum protein concentrations during, in intact and T-cell deprived mice. I. The acute phase proteins, C3 and serum amyloid P-component (SAP), 249
Serum protein concentrations during *Schistosoma mansoni* infection in intact and T-cell deprived mice. I. The acute phase proteins, C3 and serum amyloid P-component (SAP), 249
Sheep red cells (SRC), the effect of ethidium bromide on the response of spleen explants of primed rabbits to, 165
Sodium azide inhibition of complement-mediated functions, 53
Staphylococcus aureus, protein A of, is mitogenic for IgG-bearing, but also for a subpopulation of IgM- and/or IgD-bearing human lymphocytes, 417
Streptococcus mutans, comparative opsonic activity for, in oral fluids, and phagocytic activity of blood, crevicular, and salivary polymorphonuclear leucocytes in rhesus monkeys, 101
Suppressor cells, the induction of, in mixed leucocyte cultures and in mixed leucocyte–non-lymphoid cell cultures, 11

T and B lymphoma cell lines, hybridization between, 57
T antigen and T agglutinin in inbred rats, characterization of the, 219
T-cell mediated, limited specific, cytolysis in the absence of extracellular Ca^{2+}, 109
T-cell response, enhancement of, to PWM by neuraminidase-treated non-T cells, 203
T-cell responses, suppression of, to histocompatibility antigens by BCG pre-treatment, 263
T-cell subpopulations, regulatory, in pregnancy. II. Evidence for suppressive activity of the late phase of MLR, 239
T cells, development of suppressor, in mice heavily infected with mycobacteria, 367
Taenia taeniaeformis, a mechanism to account for mouse strain variation in resistance to the larval cestode, 479
Testosterone-propionate-treated chicken, the development of unusual B-cell functions in the, 29
Testosterone propionate, the B-cell development independent of the bursa of Fabricius but dependent upon the thymus in chickens treated with, 37
Thy-1 antigen: selective association in lymphoid organs with the vascular basement membrane involved in lymphocyte recirculation, 85
Thy-1 antigen, hidden, in a subpopulation of mouse bone marrow cells, 375
Tumour resistance, dissociation between, and delayed-type hypersensitivity to tumour-associated antigens in the mouse, 1
Tumours, solid, quantitative studies of natural immunity to, in rats. Persistence of natural immunity throughout reproductive life, and absence of suppressor cells in infant rats, 187

Xenoantiserum to human C3 receptors: its preparation and effect on the C3b and C3d receptors of tonsil cells and the C3b receptors of erythrocytes and neutrophils, 75